

# Emission lines of Fe XI in the 257–407 Å wavelength region observed in solar spectra from EIS/Hinode and SERTS

F. P. Keenan<sup>\*,1</sup> R. O. Milligan,<sup>2</sup> D. B. Jess,<sup>1</sup> K. M. Aggarwal,<sup>1</sup> M. Mathioudakis,<sup>1</sup>  
R. J. Thomas,<sup>2</sup> J. W. Brosius<sup>2,3</sup> and J. M. Davila<sup>2</sup>

<sup>1</sup>*Astrophysics Research Centre, School of Mathematics and Physics, Queen's University, Belfast BT7 1NN*

<sup>2</sup>*Laboratory for Solar Physics, Code 671, Heliophysics Science Division, NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA*

<sup>3</sup>*Department of Physics, The Catholic University of America, Washington, DC 20064, USA*

Received 2009 xxxx

## ABSTRACT

Theoretical emission-line ratios involving Fe XI transitions in the 257–407 Å wavelength range are derived using fully relativistic calculations of radiative rates and electron impact excitation cross sections. These are subsequently compared with both long wavelength channel Extreme-Ultraviolet Imaging Spectrometer (EIS) spectra from the *Hinode* satellite (covering 245–291 Å), and first-order observations ( $\sim 235$ –449 Å) obtained by the Solar Extreme-ultraviolet Research Telescope and Spectrograph (SERTS). The 266.39, 266.60 and 276.36 Å lines of Fe XI are detected in two EIS spectra, confirming earlier identifications of these features, and 276.36 Å is found to provide an electron density ( $N_e$ ) diagnostic when ratioed against the 257.55 Å transition. Agreement between theory and observation is found to be generally good for the SERTS data sets, with discrepancies normally being due to known line blends, while the 257.55 Å feature is detected for the first time in SERTS spectra. The most useful Fe XI electron density diagnostic is found to be the 308.54/352.67 intensity ratio, which varies by a factor of 8.4 between  $N_e = 10^8$  and  $10^{11}$  cm<sup>-3</sup>, while showing little temperature sensitivity. However, the 349.04/352.67 ratio potentially provides a superior diagnostic, as it involves lines which are closer in wavelength, and varies by a factor of 14.7 between  $N_e = 10^8$  and  $10^{11}$  cm<sup>-3</sup>. Unfortunately, the 349.04 Å line is relatively weak, and also blended with the second-order Fe X 174.52 Å feature, unless the first-order instrument response is enhanced.

**Key words:** atomic data – Sun: activity – Sun: corona – Sun: ultraviolet.

## 1 INTRODUCTION

Emission lines arising from transitions in sulphur-like Fe XI have been detected in the spectra of a range of astrophysical sources, including the Sun (Dere 1978) and other cool stars (Laming & Drake 1999), as well as active galaxies (Puchnarewicz et al. 1998). The usefulness of these lines as electron density diagnostics for the emitting plasma was first shown by Kastner & Mason (1978), who employed the atomic physics calculations of Mason (1975) to generate theoretical emission-line ratios for this ion. Since then, several authors have calculated atomic data and theoretical line ratios for Fe XI, particularly for applications to solar spectra (see Bhatia et al. 2002 and references therein).

Most work on the solar spectrum of Fe XI has fo-

cused on the  $\sim 180$ –200 Å wavelength range, containing  $3s^23p^4$ – $3s^23p^33d$  transitions (see, for example, Keenan et al. 2005). This is understandable, as this wavelength region contains numerous strong Fe XI lines (including the most intense in the solar spectrum), many of which provide good electron density diagnostics. However, the  $\sim 257$ –407 Å region also contains several Fe XI lines, primarily arising from  $3s^23p^4$ – $3s3p^5$  transitions. In this paper we use the most recent atomic physics calculations for Fe XI to analyse solar spectra covering this wavelength region. Specifically, we investigate observational data sets from two missions, namely the Extreme-Ultraviolet Imaging Spectrometer (EIS) on board the *Hinode* satellite, and the Solar Extreme-ultraviolet Research Telescope and Spectrograph (SERTS) sounding rocket experiments. The EIS long wavelength channel spectral region is 245–291 Å (Culhane et al. 2007), compared to  $\sim 235$ –449 Å for the SERTS first-order

\* E-mail: F.Keenan@qub.ac.uk

bandpass (Neupert et al. 1992), and both have spectral resolutions of about  $0.06 \text{ \AA}$  (FWHM). Although SERTS has a larger wavelength coverage, in principle allowing more lines from the same species to be detected, it has a limited dynamic range as most spectra have been recorded on photographic film. By contrast, EIS spectra are obtained with CCDs, and hence can reliably detect weaker lines. The combination of wavelength coverage and dynamic range provided by SERTS and EIS allows us to undertake a detailed assessment of the Fe XI emission-line spectrum in the 257–407  $\text{\AA}$  wavelength region.

## 2 THEORETICAL LINE RATIOS

The model ion adopted for Fe XI has been discussed in detail by Keenan et al. (2005). Briefly, it consisted of the 24 energetically lowest LS states, yielding a total of 48 fine-structure levels. Experimental energy levels, where available, were obtained from Shirai et al. (1990) and Jupén et al. (1993), with the theoretical results of Aggarwal & Keenan (2003a) being used for the remainder. Electron impact rates were taken from Aggarwal & Keenan (2003b), while for Einstein A-coefficients the data of Aggarwal & Keenan (2003a) were employed. Finally, under solar plasma conditions proton impact excitation is only important for transitions within the  $3s^23p^4 \text{ } ^3P$  ground term, and we have used the calculations of Landman (1980) for this atomic process.

Using the model ion outlined above, in conjunction with an updated version of the statistical equilibrium code of Dufton (1977), relative Fe XI emission-line strengths were calculated as a function of both electron temperature ( $T_e$ ) and density ( $N_e$ ). Details of the procedures involved and approximations made may be found in Dufton (1977) and Dufton et al. (1978). Given errors in the adopted atomic data of typically  $\pm 10$  per cent (see the references above), we would expect the theoretical ratios to be accurate to better than  $\pm 20$  per cent.

## 3 OBSERVATIONAL DATA

The solar spectra analysed in the present paper are from two instruments, namely the EIS on board the *Hinode* satellite, and several SERTS rocket flights. Our EIS data sets are those of an active region and limb area originally presented and discussed by Brown et al. (2008), to which the reader is referred for details of the EIS instrument and the data reduction and flux calibration procedures. These two features, denoted AR1 and Limb by Brown et al., show the strongest Fe XI emission lines of the solar regions considered by these authors. The spectra are from the EIS long wavelength channel and cover 245–291  $\text{\AA}$  at a resolution of about 55 m $\text{\AA}$  (FWHM).

Our SERTS spectra are those of several quiet and active regions, plus an off-limb area, obtained during rocket flights in 1989 (denoted SERTS89), 1991 (SERTS91), 1993 (SERTS93) and 1997 (SERTS97). Details of these observations, including their reduction and flux calibration, may be found in Thomas & Neupert (1994) for SERTS89, Brosius et al. (1996) for SERTS91 and SERTS93, and Brosius et al. (2000) for SERTS97. Most of these data sets

cover the 235–449  $\text{\AA}$  (SERTS89) or 231–445  $\text{\AA}$  (SERTS91 and SERTS93) wavelength regions in first-order, at resolutions of between 50–80 m $\text{\AA}$ , with spectra recorded on photographic film. However, the intensity calibrations for the SERTS91 and SERTS93 spectra are highly uncertain for wavelengths shorter than 274  $\text{\AA}$ . By contrast, the SERTS97 spectrum spans 299–353  $\text{\AA}$  in first-order, at a resolution of 115 m $\text{\AA}$ , and was recorded on a CCD. We note that the version of SERTS flown in 1989 carried a standard gold-coated toroidal diffraction grating, while those in 1991, 1993 and 1997 incorporated a multilayered-coated grating which enhanced the instrument sensitivity in the first-order wavelength range, by factors of up to 9. In addition, the absolute intensity calibration of the SERTS89 spectrum has been re-evaluated since the publication of Thomas & Neupert, as a result of which emission-line intensities are uniformly a factor of 1.24 larger than those quoted in that paper (see, for example, Keenan et al. 2008 and references therein).

## 4 RESULTS AND DISCUSSION

We have searched for Fe XI emission lines in the EIS and SERTS spectra using the detections of Brown et al. (2008) and Thomas & Neupert (1994), supplemented with those from other sources, including the National Institute of Standards and Technology (NIST) data base,<sup>1</sup> the latest version (V6.0) of the CHIANTI data base (Dere et al. 1997, 2009), and the Atomic Line List of van Hoof.<sup>2</sup> In Table 1 we list the Fe XI transitions identified in the data sets, along with their measured wavelengths. We also provide a note on each line, including whether it is detected in both EIS and SERTS spectra, and if there are possible blending features.


Intensities and line widths (FWHM) of the Fe XI features are given in Tables 2 and 3–9 for the EIS and SERTS data sets, respectively, along with the associated  $1\sigma$  errors. In the case of EIS, we have remeasured the spectra considered by Brown et al. (2008) using a standard least-squares fitting routine in SSIDL assuming Gaussian profiles and a constant background. For SERTS the measurements were made with modified versions of the Gaussian fitting routines originally employed by Thomas & Neupert (1994), as discussed by Keenan et al. (2007). As a consequence, the EIS and SERTS line intensities, FWHM values and their errors listed in Tables 2–9 are somewhat different from those originally reported in Brown et al., Thomas & Neupert and Brosius et al. (1996, 2000). However, in most directly comparable cases, the measured intensity and width values usually differ only slightly from those previously obtained. An exception is the 349.04  $\text{\AA}$  line in the SERTS spectra, where the features around 349  $\text{\AA}$  have been fitted with a triple Gaussian as opposed to the double Gaussian adopted by Thomas & Neupert and Brosius et al. This was to allow for the presence of the weak Fe X 349.30  $\text{\AA}$  transition, instead of only Mg VI 349.16  $\text{\AA}$ .

<sup>1</sup> <http://physics.nist.gov/PhysRefData/>

<sup>2</sup> <http://www.pa.uky.edu/~peter/atomic/>

**Table 1.** Fe XI line identifications in the EIS and SERTS spectra.

| Wavelength (Å) | Transition                                     | Note                                                                             |
|----------------|------------------------------------------------|----------------------------------------------------------------------------------|
| 257.55         | $3s^2 3p^4 \ ^3P_2 - 3s^2 3p^3(^4S)3d \ ^5D_3$ | Detected by both EIS and SERTS.                                                  |
| 257.77         | $3s^2 3p^4 \ ^3P_2 - 3s^2 3p^3(^4S)3d \ ^5D_2$ | Detected by EIS.                                                                 |
| 266.39         | $3s^2 3p^4 \ ^3P_1 - 3s^2 3p^3(^4S)3d \ ^5D_1$ | Detected by EIS.                                                                 |
| 266.60         | $3s^2 3p^4 \ ^3P_1 - 3s^2 3p^3(^4S)3d \ ^5D_0$ | Detected by EIS.                                                                 |
| 276.36         | $3s^2 3p^4 \ ^3P_2 - 3s3p^5 \ ^1P_1$           | Detected by EIS.                                                                 |
| 308.54         | $3s^2 3p^4 \ ^1D_2 - 3s3p^5 \ ^1P_1$           | Detected by SERTS. Blended with Fe VI (Brosius et al. 1998).                     |
| 341.11         | $3s^2 3p^4 \ ^3P_2 - 3s3p^5 \ ^3P_1$           | Detected by SERTS.                                                               |
| 349.04         | $3s^2 3p^4 \ ^3P_1 - 3s3p^5 \ ^3P_0$           | Detected by SERTS. Blended with 2nd-order Fe X 174.52 Å (Thomas & Neupert 1994). |
| 352.67         | $3s^2 3p^4 \ ^3P_2 - 3s3p^5 \ ^3P_2$           | Detected by SERTS.                                                               |
| 356.53         | $3s^2 3p^4 \ ^3P_1 - 3s3p^5 \ ^3P_1$           | Detected by SERTS.                                                               |
| 358.67         | $3s^2 3p^4 \ ^3P_0 - 3s3p^5 \ ^3P_1$           | Detected by SERTS. Blended with several lines (Young et al. 1998).               |
| 369.16         | $3s^2 3p^4 \ ^3P_1 - 3s3p^5 \ ^3P_2$           | Detected by SERTS.                                                               |
| 406.79         | $3s^2 3p^4 \ ^1D_2 - 3s3p^5 \ ^3P_2$           | Detected by SERTS. Identified in spectrum by Brickhouse et al. (1995).           |

**Figure 1.** Portions of the EIS active region and limb spectra containing the Fe XI 257.55, 257.77, 266.39, 266.60 and 276.36 Å emission lines. The profile fits to these features are shown by dashed lines.**Table 2.** Fe XI line intensities and widths from the EIS spectra.

| Feature <sup>a</sup> | Wavelength (Å) | Intensity (erg cm <sup>-2</sup> s <sup>-1</sup> sr <sup>-1</sup> ) | Line width (mÅ) |
|----------------------|----------------|--------------------------------------------------------------------|-----------------|
| Active region        | 257.55         | 335.0 ± 31.1                                                       | 73 ± 5          |
|                      | 257.77         | 167.0 ± 24.2                                                       | 55 ± 8          |
|                      | 266.39         | 17.0 ± 3.4                                                         | 70 ± 4          |
|                      | 266.60         | 107.0 ± 9.1                                                        | 90 ± 4          |
|                      | 276.36         | 16.0 ± 7.7                                                         | 82 ± 10         |
| Limb                 | 257.55         | 278 ± 23.9                                                         | 77 ± 5          |
|                      | 257.77         | 133.0 ± 14.2                                                       | 59 ± 9          |
|                      | 266.39         | 19.0 ± 3.3                                                         | 96 ± 14         |
|                      | 266.60         | 45.0 ± 6.8                                                         | 100 ± 9         |
|                      | 276.36         | 6.0 ± 2.5                                                          | 85 ± 10         |

<sup>a</sup>The active region and limb features are those denoted AR1 and Limb, respectively, by Brown et al. (2008).**Table 3.** Fe XI line intensities and widths from the SERTS 1989 active region spectrum (SERTS89-AR).

| Wavelength (Å)      | Intensity (erg cm <sup>-2</sup> s <sup>-1</sup> sr <sup>-1</sup> ) | Line width (mÅ) |
|---------------------|--------------------------------------------------------------------|-----------------|
| 257.55              | 35.5 ± 15.1                                                        | 96 ± 10         |
| 308.54 <sup>a</sup> | 106.5 ± 20.3                                                       | 122 ± 12        |
| 341.11              | 46.1 ± 6.9                                                         | 93 ± 8          |
| 349.04              | 26.2 ± 11.9                                                        | 105 ± 39        |
| 352.67              | 161.6 ± 18.8                                                       | 95 ± 6          |
| 356.53              | 22.1 ± 4.8                                                         | 92 ± 17         |
| 358.67              | 88.7 ± 11.4                                                        | 144 ± 9         |
| 369.16              | 47.1 ± 6.3                                                         | 102 ± 5         |
| 406.79              | 3.6 ± 1.4                                                          | 57 ± 13         |

<sup>a</sup>Line measured at wavelength of 308.58 Å in this spectrum (see Section 4.2 for details).

**Table 4.** Fe XI line intensities and widths from the SERTS 1991 active region spectrum (SERTS91-AR).

| Wavelength<br>(Å) | Intensity<br>(erg cm <sup>-2</sup> s <sup>-1</sup> sr <sup>-1</sup> ) | Line width<br>(mÅ) |
|-------------------|-----------------------------------------------------------------------|--------------------|
| 308.54            | 5.9 ± 1.7                                                             | 40 ± 10            |
| 341.11            | 20.3 ± 3.7                                                            | 56 ± 12            |
| 352.67            | 36.8 ± 5.6                                                            | 57 ± 7             |
| 358.67            | 19.4 ± 4.9                                                            | 64 ± 20            |
| 369.16            | 13.4 ± 3.6                                                            | 69 ± 10            |

**Table 5.** Fe XI line intensities and widths from the SERTS 1991 quiet Sun spectrum (SERTS91-QS).

| Wavelength<br>(Å) | Intensity<br>(erg cm <sup>-2</sup> s <sup>-1</sup> sr <sup>-1</sup> ) | Line width<br>(mÅ) |
|-------------------|-----------------------------------------------------------------------|--------------------|
| 308.54            | 19.2 ± 5.7                                                            | 38 ± 5             |
| 341.11            | 58.6 ± 10.3                                                           | 81 ± 14            |
| 352.67            | 112.6 ± 16.1                                                          | 58 ± 10            |
| 356.53            | 17.2 ± 7.8                                                            | 45 ± 15            |
| 358.67            | 14.2 ± 8.1                                                            | 41 ± 16            |
| 369.16            | 44.1 ± 12.3                                                           | 70 ± 22            |

**Table 6.** Fe XI line intensities and widths from the SERTS 1991 off-limb spectrum (SERTS91-OL).

| Wavelength<br>(Å) | Intensity<br>(erg cm <sup>-2</sup> s <sup>-1</sup> sr <sup>-1</sup> ) | Line width<br>(mÅ) |
|-------------------|-----------------------------------------------------------------------|--------------------|
| 308.54            | 42.8 ± 13.1                                                           | 43 ± 10            |
| 341.11            | 99.7 ± 15.1                                                           | 94 ± 12            |
| 352.67            | 240.8 ± 29.1                                                          | 72 ± 8             |
| 358.67            | 36.3 ± 18.2                                                           | 37 ± 10            |
| 369.16            | 59.9 ± 8.3                                                            | 109 ± 30           |

**Table 7.** Fe XI line intensities and widths from the SERTS 1993 active region spectrum (SERTS93-AR).

| Wavelength<br>(Å) | Intensity<br>(erg cm <sup>-2</sup> s <sup>-1</sup> sr <sup>-1</sup> ) | Line width<br>(mÅ) |
|-------------------|-----------------------------------------------------------------------|--------------------|
| 308.54            | 28.7 ± 11.6                                                           | 45 ± 15            |
| 341.11            | 71.1 ± 10.1                                                           | 62 ± 9             |
| 352.67            | 178.1 ± 20.5                                                          | 57 ± 9             |
| 356.53            | 19.3 ± 8.5                                                            | 42 ± 25            |
| 358.67            | 61.1 ± 11.5                                                           | 100 ± 17           |
| 369.16            | 59.8 ± 13.4                                                           | 103 ± 22           |

**Table 8.** Fe XI line intensities and widths from the SERTS 1993 quiet Sun spectrum (SERTS93-QS).

| Wavelength<br>(Å) | Intensity<br>(erg cm <sup>-2</sup> s <sup>-1</sup> sr <sup>-1</sup> ) | Line width<br>(mÅ) |
|-------------------|-----------------------------------------------------------------------|--------------------|
| 308.54            | 6.6 ± 3.2                                                             | 32 ± 12            |
| 341.11            | 13.5 ± 2.8                                                            | 50 ± 10            |
| 352.67            | 48.3 ± 6.5                                                            | 63 ± 9             |
| 358.67            | 10.3 ± 2.9                                                            | 83 ± 16            |
| 369.16            | 13.2 ± 3.4                                                            | 115 ± 23           |

**Table 9.** Fe XI line intensities and widths from the SERTS 1997 active region spectrum (SERTS97-AR).

| Wavelength<br>(Å) | Intensity<br>(erg cm <sup>-2</sup> s <sup>-1</sup> sr <sup>-1</sup> ) | Line width<br>(mÅ) |
|-------------------|-----------------------------------------------------------------------|--------------------|
| 308.54            | 33.9 ± 9.8                                                            | 142 ± 8            |
| 341.11            | 37.8 ± 6.7                                                            | 149 ± 9            |
| 349.04            | 12.5 ± 4.2                                                            | 103 ± 9            |
| 352.67            | 136.1 ± 22.1                                                          | 135 ± 9            |

#### 4.1 EIS spectra


Although CHIANTI lists 23 transitions of Fe XI in the EIS long wavelength channel (245–291 Å), most of these are either too weak to be detected or are blended with much stronger emission features from other species. As a result, we only have measurements for 5 lines, which are summarised in Table 2. Portions of the EIS active region and limb spectra containing these lines are shown in Figure 1. In Table 10 we list the resultant observed line ratios (and their 1 $\sigma$  errors), along with the theoretical results from the present calculations. Also given are the theoretical predictions from the CHIANTI V6.0 data base, which employs the electron excitation rates of Gupta & Tayal (1999) and Bhatia & Doschek (1996). The 266.39/257.55 and 266.60/257.55 ratios are relatively insensitive to changes in the electron density over the range  $N_e = 10^9$ – $10^{11}$  cm<sup>-3</sup>, varying by only 13 and 8 per cent, respectively. However, 276.36/257.55 is  $N_e$ -sensitive, as may be seen in Figure 2. Hence we have determined values of  $N_e$  for the active region and limb using the 200.03/202.04 and 203.17/202.04 line intensity ratios in Fe XIII, which is formed at a similar temperature to Fe XI in ionisation equilibrium ( $T_e = 10^{6.2}$  K compared to  $10^{6.1}$  K for Fe XI; Bryans et al. 2009). For the active region, we measured 200.03/202.04 = 0.31 and 203.17/202.04 = 0.16, both of which yield  $N_e = 10^{9.2}$  cm<sup>-3</sup> from the calculations of Keenan et al. (2007). Similarly, for the limb spectrum the observed ratios are 200.03/202.04 = 0.47 and 203.17/202.04 = 0.24, which both indicate  $N_e = 10^{9.5}$  cm<sup>-3</sup>. In Table 10 we therefore list the theoretical ratios generated at the temperature of maximum Fe XI fractional abundance in ionisation equilibrium and at densities of  $10^{9.2}$  and  $10^{9.5}$  cm<sup>-3</sup> for the active region and limb area, respectively.

The 266.39 Å line was tentatively identified by Träbert (1998) in the solar flare spectra of Dere (1978), and the reasonable agreement between theory and observation for the 266.39/257.55 ratio in Table 10 (especially for the CHIANTI calculations) supports this identification. We note that CHIANTI lists the  $3s^2 3p^4 \ ^3P_1$ – $3s^2 3p^3 (^4S) 3d \ ^5D_1$  transition at 273.41 Å, but no emission feature is detected at this wavelength in the EIS spectra. In addition, Brown et al. (2008) identify the 266.39 Å feature as an Fe XVII transition. However, the fact that the measured 266.39/257.55 ratios are actually somewhat smaller than theory indicates that Fe XVII probably makes a negligible contribution to the 266.39 Å line intensity.

Träbert (1998) also tentatively identified the 266.60 Å line of Fe XI in the Dere (1978) spectra, but in this case the experimental 266.60/257.55 ratios in Table 10 are larger than the theoretical values, at least for the active region. This suggests that the 266.60 Å feature is blended, which

**Table 10.** Fe XI line ratios from the EIS spectra.

| Feature       | Line ratio    | Observed          | Present theory <sup>a</sup> | CHIANTI theory <sup>a</sup> |
|---------------|---------------|-------------------|-----------------------------|-----------------------------|
| Active region | 257.77/257.55 | $0.50 \pm 0.09$   | 0.62                        | 0.52                        |
| Active region | 266.39/257.55 | $0.051 \pm 0.011$ | 0.12                        | 0.083                       |
| Active region | 266.60/257.55 | $0.32 \pm 0.04$   | 0.11                        | 0.069                       |
| Active region | 276.36/257.55 | $0.048 \pm 0.024$ | 0.032                       | 0.038                       |
| Limb          | 257.77/257.55 | $0.48 \pm 0.07$   | 0.61                        | 0.54                        |
| Limb          | 266.39/257.55 | $0.068 \pm 0.013$ | 0.12                        | 0.084                       |
| Limb          | 266.60/257.55 | $0.16 \pm 0.03$   | 0.11                        | 0.072                       |
| Limb          | 276.36/257.55 | $0.022 \pm 0.009$ | 0.041                       | 0.051                       |

<sup>a</sup>Present theoretical ratios and those from CHIANTI calculated at  $T_e = 10^{6.1}$  K and  $N_e = 10^{9.2} \text{ cm}^{-3}$  (active region) or  $N_e = 10^{9.5} \text{ cm}^{-3}$  (limb).**Figure 2.** The theoretical Fe XI emission-line intensity ratio  $I(276.36 \text{ Å})/I(257.55 \text{ Å})$ , where  $I$  is in energy units, plotted as a function of logarithmic electron density ( $N_e$  in  $\text{cm}^{-3}$ ) at the temperature of maximum Fe XI fractional abundance in ionization equilibrium,  $T_e = 10^{6.1}$  K (Bryans et al. 2009), plus  $\pm 0.2$  dex about this value.

is supported by the relatively large width found for the line. However, an inspection of CHIANTI and other line lists reveals no likely blending candidates. We note that CHIANTI lists a wavelength of  $273.57 \text{ Å}$  for the  $3s^2 3p^4 \text{ } ^3\text{P}_1 - 3s^2 3p^3 ({}^4\text{S}) 3d \text{ } ^5\text{D}_0$  transition, which unfortunately would be blended with Fe XIV  $273.55 \text{ Å}$ . However, the experimental Fe XIV  $273.55/274.20$  ratios in the EIS spectra are in good agreement with the CHIANTI predictions, indicating that Fe XI does not make a contribution to the  $273.55 \text{ Å}$  line intensity. We also note that Brown et al. (2008) detect the  $266.60 \text{ Å}$  line, but no identification is provided.

Of particular interest is the  $276.36 \text{ Å}$  line, which was unidentified in solar observations until recently detected by Brown et al. (2008) in the EIS limb spectrum. However, these authors identify the feature as Fe XII rather than Fe XI, although this is almost certainly a typographical error as the transition is correctly given as  $3s^2 3p^4 \text{ } ^3\text{P}_2 - 3s 3p^5 \text{ } ^1\text{P}_1$ . (We note in passing that Brown et al. incorrectly list the  $257.77 \text{ Å}$  line as having a  ${}^5\text{D}_4$  upper level, when it should be  ${}^5\text{D}_2$ ). An inspection of Table 10 reveals good agreement between theory and observation for  $276.36/257.55$ , confirming the identification of the  $276.36 \text{ Å}$  feature as the  $3s^2 3p^4 \text{ } ^3\text{P}_2 - 3s 3p^5 \text{ } ^1\text{P}_1$  line of Fe XI. However, further observations of the transition would be useful, as the  $276.36/257.55$  ratio potentially provides a useful EIS long wavelength channel density diagnostic for the Fe XI-emitting region of a plasma, as it varies by a factor of 6.8 between  $N_e = 10^8$  and  $10^{12} \text{ cm}^{-3}$ .

In Table 10 we also list the observed and theoretical  $257.77/257.55$  ratios, for which there is good agreement, indicating that both lines are well resolved in the EIS spectra and free from blends. To check this we have generated synthetic active region, flare and quiet Sun spectra using V6.0 of the CHIANTI data base. These reveal no significant blends for either the  $257.55$  or  $257.77 \text{ Å}$  transitions, nor indeed for the  $276.36 \text{ Å}$  feature. We note that Jupén et al. (1993) originally identified the  $257.55$  and  $257.77 \text{ Å}$  features as being the (2–3) and (2–2) components, respectively, of the  $3s^2 3p^4 \text{ } ^3\text{P}_J - 3s^2 3p^3 ({}^4\text{S}) 3d \text{ } ^5\text{D}_J$  multiplet. Subsequently, Träbert (1998) reclassified the lines as the (2–2) and (2–1) components, with (2–3) lying at  $257.26 \text{ Å}$  and hence blended with a strong Fe X line. However, the theoretical (2–1)/(2–2) ratio is 0.31 (present calculations) or 0.27 (CHIANTI), much smaller than the experimental values of around 0.50 for  $257.77/257.55$ . Hence the EIS observations do not support the Träbert classifications, but rather those of Jupén et al. A definitive answer would be provided by the unambiguous detection of the (1–2) or (2–1) components of the multiplet. Unfortunately, the (1–2) transition is predicted to be too weak to observe, with a theoretical intensity of less than 0.1 per cent that of the  $257.55 \text{ Å}$  feature. Similarly, the (2–1) transition is predicted to lie at  $264.92 \text{ Å}$  by CHIANTI, and cannot be observed as it lies between lines of Fe XIV and Fe XVI.

## 4.2 SERTS observations

The SERTS first-order wavelength range ( $\sim 235\text{--}449 \text{ Å}$ ) contains numerous Fe XI emission features, resulting in many possible line ratio combinations. Following Keenan et al. (2007), we categorise 3 types of ratio, namely:

- (i) branching ratios, where the transitions arise from a common upper level. In this instance the ratio is predicted to be constant under most plasma conditions, an exception being when significant opacity is present, which should not apply to the Fe XI lines considered here;
- (ii) ratios which are predicted to be relatively insensitive to changes in  $T_e$  and  $N_e$  over the range of plasma parameters of interest;
- (iii) ratios which are predicted to vary significantly with  $N_e$ , and hence may provide useful electron density diagnostics.

Ratios in categories (i) and (ii) are the most useful

for identifying and assessing the importance of blends, as well as investigating possible errors in the adopted atomic data, as one does not need to accurately know the plasma parameters to generate a theoretical ratio for comparison with observations. In Tables 11 and 12 we therefore list the observed line ratios for the SERTS data sets which fall into category (i) or (ii), respectively (along with the associated  $1\sigma$  errors), as well as the theoretical values both from the present calculations and CHIANTI. Once again, following Keenan et al. (2007), we have defined ‘relatively insensitive’ as being those ratios which are predicted to vary by less than  $\pm 20$  per cent when the electron density is changed by a factor of 2 (i.e.  $\pm 0.3$  dex). In fact, most of the ratios in Table 12 show much lower sensitivity to variations in  $N_e$ . For example, at  $T_e = 10^{6.1}$  K, the 341.11/352.67 ratio only changes from 0.27 at  $N_e = 10^9 \text{ cm}^{-3}$  to 0.31 at  $N_e = 10^{10} \text{ cm}^{-3}$ , while 257.55/352.67 varies from 0.13 to 0.16 over the same density interval. As noted by Keenan et al., most of the electron densities derived for SERTS active regions from species formed at similar temperatures to Fe XI are consistent with  $\log N_e = 9.4 \pm 0.3$ . This is also the case for both quiet Sun regions and the off-limb area, where diagnostics indicate  $\log N_e = 9.1 \pm 0.3$  (Brosius et al. 1996). The theoretical results in Table 12 have therefore been calculated at the temperature of maximum Fe XI fractional abundance in ionization equilibrium,  $T_e = 10^{6.1}$  K (Bryans et al. 2009), and at  $N_e = 10^{9.4} \text{ cm}^{-3}$  (for the active regions) or  $N_e = 10^{9.1} \text{ cm}^{-3}$  (quiet Sun and off-limb). However, we note that changing the adopted value of  $T_e$  by  $\pm 0.2$  dex or the density by  $\pm 0.5$  dex does not significantly alter the discussions below.

An inspection of Table 12 reveals good agreement between theory and observation for the 257.55/352.67 ratio, providing support for our identification of the 257.55 Å line. This is the first time this feature has been detected in SERTS observations, although it has previously been identified by Jupén et al. (1993) in the solar spectrum of Behring et al. (1976), as well as in EIS data (see Section 4.1).

The measured 349.04/308.54 ratio in Table 12 for the SERTS89 active region is smaller than the present theoretical result or that from CHIANTI, which indicates blending in the 308.54 Å line. Brosius et al. (1998) note that the Fe XI 308.54 Å feature is blended with an Fe VI transition, but from the CHIANTI synthetic spectra the latter is predicted to contribute less than 0.1 per cent of the total intensity. Indeed, CHIANTI indicates that there are lines of Ni XIII, Al VI and Ne III which make a larger contribution to the blend than Fe VI, but even then the summed intensity of these 3 transitions is only 5 per cent that of Fe XI. On the other hand, the 308.54 Å line width in the SERTS89 active region spectrum is much larger than in the other SERTS data sets, as may be seen from Figure 3 and Tables 3–9. Furthermore, the feature lies at 308.58 Å in the SERTS89 observations, compared to 308.54 Å in the remaining SERTS data sets and other solar spectra, such as those from the S082A instrument on board *Skylab* (Dere 1978). Both the large line width and wavelength shift are symptomatic of blending, but if such a species is present it is difficult to explain why it should affect one active region and not others. A more likely explanation is a fault with the SERTS spectrum, although an inspection of the film reveals no apparent flaws.

The experimental 349.04/308.54 ratio for the SERTS97




**Figure 3.** Plot of the SERTS 1989 (lower panel) and 1993 (upper panel) active region spectra in the 307.8–309.3 Å wavelength range. The profile fits to the Fe XI 308.54 Å feature are shown by solid lines.

active region is in good agreement with theory, indicating that both these features are reliably observed and free from blends. Thomas & Neupert (1994) point out that the Fe XI 349.04 Å line in the SERTS89 data is blended with the second-order Fe X 174.52 Å transition, which appears at 349.04 Å in first-order. However, as noted in Section 3, the version of SERTS flown in 1997 incorporated a multilayered-coated grating, which enhanced the instrument sensitivity in first-order compared to the SERTS89 observations, hence removing the effect of the blend with Fe X 174.52 Å. On the other hand, this blending is evident in the spectrum from the SERTS flight in 1995, where the instrument sensitivity was enhanced in the second-order waveband (Brosius et al. 1998), and the observed 349.04/308.54 ratio is 3.4, much larger than the theoretical value of about 0.6.

Both the experimental 341.11/356.53 and 341.11/352.67 ratios in Tables 11 and 12 are generally larger than the theoretical values, which CHIANTI indicates is due to blending of 341.11 Å with an Fe IX transition. The latter is predicted to contribute about 33 per cent to the total 341.11 Å line intensity, which is consistent with the observations. For example, for the SERTS93 active region, reducing the intensity of the 341.11 Å line by 33 per cent leads to a revised 341.11/352.67 ratio of 0.26, in good agreement with the theoretical value of 0.29.

The measured 356.53/352.67 ratios in Table 12 are all in good agreement with theory, indicating that both the 352.67 and 356.53 Å lines are well observed and free from blends. Similarly, there are no discrepancies between theory and observation for 369.16/352.67, and indeed the average of the 6 measurements in Table 11 is  $0.32 \pm 0.05$ , which compares very well with the theoretical result of 0.30, showing that the 369.16 Å transition is also free from blends. We note that the 352.67, 356.53 and 369.16 Å lines of Fe XI have also

**Table 11.** Comparison of theory and observation for SERTS line ratios involving transitions from common upper levels.

| Feature    | Line ratio    | Observed          | Present theory | CHIANTI theory |
|------------|---------------|-------------------|----------------|----------------|
| SERTS89-AR | 341.11/356.53 | $2.1 \pm 0.6$     | 2.0            | 2.0            |
| SERTS91-QS | 341.11/356.53 | $3.4 \pm 1.7$     | 2.0            | 2.0            |
| SERTS93-AR | 341.11/356.53 | $3.7 \pm 1.7$     | 2.0            | 2.0            |
| SERTS89-AR | 341.11/358.67 | $0.52 \pm 0.10$   | 1.6            | 1.6            |
| SERTS91-AR | 341.11/358.67 | $1.0 \pm 0.3$     | 1.6            | 1.6            |
| SERTS93-QS | 341.11/358.67 | $1.3 \pm 0.5$     | 1.6            | 1.6            |
| SERTS89-AR | 358.67/356.53 | $4.0 \pm 1.0$     | 1.2            | 1.3            |
| SERTS91-QS | 358.67/356.53 | $0.83 \pm 0.60$   | 1.2            | 1.3            |
| SERTS93-AR | 358.67/356.53 | $3.2 \pm 1.5$     | 1.2            | 1.3            |
| SERTS89-AR | 369.16/352.67 | $0.29 \pm 0.05$   | 0.30           | 0.30           |
| SERTS91-AR | 369.16/352.67 | $0.36 \pm 0.11$   | 0.30           | 0.30           |
| SERTS91-QS | 369.16/352.67 | $0.39 \pm 0.12$   | 0.30           | 0.30           |
| SERTS91-OL | 369.16/352.67 | $0.25 \pm 0.05$   | 0.30           | 0.30           |
| SERTS93-AR | 369.16/352.67 | $0.34 \pm 0.09$   | 0.30           | 0.30           |
| SERTS93-QS | 369.16/352.67 | $0.27 \pm 0.08$   | 0.30           | 0.30           |
| SERTS89-AR | 406.79/352.67 | $0.022 \pm 0.009$ | 0.023          | 0.027          |

**Table 12.** Comparison of theory and observation for SERTS line ratios which are only weakly  $N_e$ -dependent.

| Feature    | Line ratio    | Observed        | Present theory <sup>a</sup> | CHIANTI theory <sup>a</sup> |
|------------|---------------|-----------------|-----------------------------|-----------------------------|
| SERTS89-AR | 257.55/352.67 | $0.22 \pm 0.10$ | 0.14                        | 0.19                        |
| SERTS89-AR | 341.11/352.67 | $0.29 \pm 0.05$ | 0.29                        | 0.28                        |
| SERTS91-AR | 341.11/352.67 | $0.55 \pm 0.13$ | 0.29                        | 0.28                        |
| SERTS91-QS | 341.11/352.67 | $0.52 \pm 0.12$ | 0.28                        | 0.27                        |
| SERTS91-OL | 341.11/352.67 | $0.41 \pm 0.08$ | 0.28                        | 0.27                        |
| SERTS93-AR | 341.11/352.67 | $0.40 \pm 0.07$ | 0.29                        | 0.28                        |
| SERTS93-QS | 341.11/352.67 | $0.28 \pm 0.07$ | 0.28                        | 0.27                        |
| SERTS97-AR | 341.11/352.67 | $0.28 \pm 0.07$ | 0.29                        | 0.28                        |
| SERTS89-AR | 349.04/308.54 | $0.25 \pm 0.12$ | 0.65                        | 0.50                        |
| SERTS97-AR | 349.04/308.54 | $0.37 \pm 0.16$ | 0.65                        | 0.50                        |
| SERTS89-AR | 356.53/352.67 | $0.14 \pm 0.03$ | 0.14                        | 0.14                        |
| SERTS91-QS | 356.53/352.67 | $0.15 \pm 0.07$ | 0.14                        | 0.13                        |
| SERTS93-AR | 356.53/352.67 | $0.11 \pm 0.05$ | 0.14                        | 0.14                        |
| SERTS89-AR | 358.67/352.67 | $0.55 \pm 0.10$ | 0.18                        | 0.18                        |
| SERTS91-AR | 358.67/352.67 | $0.53 \pm 0.16$ | 0.18                        | 0.18                        |
| SERTS91-QS | 358.67/352.67 | $0.13 \pm 0.08$ | 0.17                        | 0.17                        |
| SERTS91-OL | 358.67/352.67 | $0.15 \pm 0.08$ | 0.17                        | 0.17                        |
| SERTS93-AR | 358.67/352.67 | $0.34 \pm 0.08$ | 0.18                        | 0.18                        |
| SERTS93-QS | 358.67/352.67 | $0.21 \pm 0.07$ | 0.17                        | 0.17                        |

<sup>a</sup>Present theoretical ratios and those from CHIANTI calculated at  $T_e = 10^{6.1}$  K and  $N_e = 10^{9.4} \text{ cm}^{-3}$  (for active regions) or  $N_e = 10^{9.1} \text{ cm}^{-3}$  (quiet Sun and off-limb regions).

been recently observed by the EUNIS solar instrument (Brosius et al. 2008), and show good agreement between theory and experiment, with measured ratios of  $356.53/352.67 = 0.16 \pm 0.03$  and  $369.16/352.67 = 0.30 \pm 0.05$ , compared to the theoretical estimates of 0.14 and 0.30, respectively.

Young et al. (1998) note that the 358.67 Å line of Fe XI is blended with several transitions of Si XI, Ne IV and Fe XIV. However, the observed 358.67/356.53 and 358.67/352.67 ratios in Tables 11 and 12 are only much larger than the theoretical values for the active region spectra, with measurements for the quiet Sun and off-limb regions being in good agreement with theory. This implies that blending of 358.67 Å is only significant in active regions. By contrast, the synthetic active region and quiet Sun spectra generated with CHIANTI predict that Fe XI should make a smaller contribu-

tion to the total 358.67 Å line intensity in the latter (less than 20 per cent compared to about 35 per cent in an active region), and hence that blending of the Fe XI feature should in fact be more severe in quiet Sun spectra. However, Fe XI observations from other solar missions support the present findings. For example, for a quiet solar region observed with the CDS instrument on the SOHO satellite, Landi et al. (2002) measured Fe XI ratios of  $358.67/356.53 = 1.1 \pm 0.1$  and  $358.67/352.67 = 0.14 \pm 0.02$ , in excellent agreement with the theoretical values of 1.2 and 0.17, respectively. Better agreement between theory and observation for the quiet Sun compared to active regions is indicative of blending with lines formed at higher temperatures than Fe XI. Our results would therefore suggest that Si XI and Fe XIV (both of which are formed at slightly higher temperatures than Fe XI) make




**Figure 4.** The theoretical Fe XI emission-line intensity ratio  $I(308.54 \text{ \AA})/I(352.67 \text{ \AA})$ , where  $I$  is in energy units, plotted as a function of logarithmic electron density ( $N_e$  in  $\text{cm}^{-3}$ ) at the temperature of maximum Fe XI fractional abundance in ionization equilibrium,  $T_e = 10^{6.1} \text{ K}$  (Bryans et al. 2009), plus  $\pm 0.2$  dex about this value.




**Figure 5.** The theoretical Fe XI emission-line intensity ratio  $I(349.04 \text{ \AA})/I(352.67 \text{ \AA})$ , where  $I$  is in energy units, plotted as a function of logarithmic electron density ( $N_e$  in  $\text{cm}^{-3}$ ) at the temperature of maximum Fe XI fractional abundance in ionization equilibrium,  $T_e = 10^{6.1} \text{ K}$  (Bryans et al. 2009), plus  $\pm 0.2$  dex about this value.

a larger contribution to the  $358.67 \text{ \AA}$  line intensity than currently predicted by CHIANTI, while Ne IV makes a smaller contribution.

Finally, the observed  $406.79/352.67$  ratio in Table 11 is in very good agreement with the present calculations and also those from CHIANTI, supporting the identification of the  $406.79 \text{ \AA}$  line in the SERTS89 active region spectrum by Brickhouse et al. (1995).

### 4.3 Electron density diagnostics from SERTS spectra

In Figures 4 and 5 we plot the electron density diagnostics  $308.54/352.67$  and  $349.04/352.67$ , respectively, while in Table 13 we list the observed ratios (and the associated  $1\sigma$  errors) along with the values of  $N_e$  derived using the theoretical results at the temperature of maximum Fe XI fractional abundance in ionisation equilibrium,  $T_e = 10^{6.1} \text{ K}$  (Bryans et al. 2009). However, we note that changing  $T_e$  by  $\pm 0.2$  dex would lead to a variation in the derived values of  $N_e$  of at most  $\pm 0.1$  dex. Also listed in the table is the factor by which the relevant ratio is predicted to vary between  $N_e = 10^8$  and  $10^{11} \text{ cm}^{-3}$ .

An inspection of Table 13 reveals that the densities de-

rived from  $308.54/352.67$  are generally in very good agreement with those estimated for the SERTS solar features from line ratios in species formed at similar temperatures to Fe XI, which as noted in Section 4.2 average about  $\log N_e = 9.4 \pm 0.3$  for active regions and  $\log N_e = 9.1 \pm 0.3$  for quiet Sun and off-limb areas. Hence we can state that, in general, the  $308.54/352.67$  ratio appears to provide a good  $N_e$ -diagnostic. The exception is the SERTS89 active region, where the derived density is more than an order of magnitude larger than expected. However, as already discussed in Section 4.2, there are problems with the  $308.54 \text{ \AA}$  feature in the SERTS89 active region observations which make its measurement uncertain.

The electron density derived for the SERTS97 active region from the  $349.04/352.67$  ratio is in good agreement with those found from other species, but in the case of the SERTS89 active region the density is much higher than expected. However, as noted by Thomas & Neupert (1994), the  $349.04 \text{ \AA}$  line in the SERTS89 observations is blended with the second-order Fe X  $174.52 \text{ \AA}$  feature. We can estimate its contribution using the nearby Fe X  $177.24 \text{ \AA}$  transition, which appears in first-order at  $354.48 \text{ \AA}$ . The first-order intensity of this line is  $3.6 \text{ erg cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$ , and the  $174.52 \text{ \AA}$  feature is predicted to be a factor of 1.8 stronger (Keenan et al. 2008), yielding an intensity of  $6.5 \text{ erg cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$ . This in turn gives a corrected intensity for Fe XI  $349.04 \text{ \AA}$  of  $19.7 \text{ erg cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$ , and revised experimental  $349.04/352.67$  ratio of 0.12. From Figure 5, this implies  $N_e = 10^{9.5} \text{ cm}^{-3}$ , in good agreement with other diagnostics.

Although the  $308.54/352.67$  ratio is probably the most useful Fe XI density diagnostic currently available in the  $257\text{--}407 \text{ \AA}$  wavelength range,  $349.04/352.67$  is superior in many respects. The lines are closer together, reducing the effects of possible errors in the instrument intensity calibration, and the ratio varies by a larger factor over the density range  $N_e = 10^8\text{--}10^{11} \text{ cm}^{-3}$  (14.7 c.f. 8.4 for  $308.54/352.67$ ). However, the  $349.04 \text{ \AA}$  feature is relatively weak, and is also blended with Fe X  $174.52 \text{ \AA}$ , unless the first-order instrument response is enhanced, as for the SERTS97 observations.

### ACKNOWLEDGMENTS

DBJ and KMA acknowledge financial support from STFC and EPSRC, respectively. ROM acknowledges support from the NASA Postdoctoral Program at the Goddard Space Flight Center, administered by Oak Ridge Associated Universities through a contract with NASA. *Hinode* is a Japanese mission developed and launched by ISAS/JAXA, collaborating with NAOJ as a domestic partner, NASA and STFC as international partners. Scientific operation of the *Hinode* mission is conducted by the *Hinode* science team organised at ISAS/JAXA. Support for the postlaunch operation is provided by JAXA and NAOJ, STFC, NASA, ESA and NSC (Norway). The SERTS rocket programme is supported by RTOP grants from the Solar Physics Office of NASA's Space Physics Division. JWB acknowledges additional NASA support under grant NAG5-13321. FPK is grateful to AWE Aldermaston for the award of a William Penney Fellowship. The authors thank Peter van Hoof for the use of his Atomic Line List. CHIANTI is a collaborative project involving the Naval Research Laboratory (USA),

**Table 13.** Electron density diagnostic line ratios from the SERTS spectra.

| Feature    | Line ratio    | Observed          | $\log N_e^a$           | Ratio variation <sup>b</sup> |
|------------|---------------|-------------------|------------------------|------------------------------|
| SERTS89-AR | 308.54/352.67 | $0.66 \pm 0.15$   | $10.7^{+0.5}_{-0.3}$   | 8.4                          |
| SERTS91-AR | 308.54/352.67 | $0.16 \pm 0.05$   | $9.4^{+0.2}_{-0.5}$    | 8.4                          |
| SERTS91-QS | 308.54/352.67 | $0.17 \pm 0.06$   | $9.4^{+0.5}_{-0.5}$    | 8.4                          |
| SERTS91-OL | 308.54/352.67 | $0.18 \pm 0.06$   | $9.5^{+0.2}_{-0.5}$    | 8.4                          |
| SERTS93-AR | 308.54/352.67 | $0.16 \pm 0.07$   | $9.4^{+0.3}_{-1.3}$    | 8.4                          |
| SERTS93-QS | 308.54/352.67 | $0.14 \pm 0.07$   | $9.2^{+0.4}_{-\infty}$ | 8.4                          |
| SERTS97-AR | 308.54/352.67 | $0.25 \pm 0.08$   | $9.8^{+0.2}_{-0.4}$    | 8.4                          |
| SERTS89-AR | 349.04/352.67 | $0.16 \pm 0.08$   | $9.9^{+0.7}_{-0.3}$    | 14.7                         |
| SERTS97-AR | 349.04/352.67 | $0.092 \pm 0.034$ | $9.3^{+0.3}_{-0.4}$    | 14.7                         |

<sup>a</sup>Determined from present line ratio calculations at  $T_e = 10^{6.1}$  K;  $N_e$  in  $\text{cm}^{-3}$ .

<sup>b</sup>Factor by which the theoretical line ratio varies between  $N_e = 10^8$  and  $10^{11} \text{ cm}^{-3}$ .

Rutherford Appleton Laboratory (UK), and the Universities of Florence (Italy) and Cambridge (UK). We are very grateful to the referee, Peter Young, for his comments on an earlier version of the paper, in particular regarding the analysis of the EIS spectra.

## REFERENCES

- Aggarwal K.M., Keenan F.P., 2003a, MNRAS, 338, 412  
 Aggarwal K.M., Keenan F.P., 2003b, A&A, 399, 799  
 Behring W.E., Cohen L., Feldman U., Doschek G.A., 1976, ApJ, 203, 521  
 Bhatia A.K., Doschek G.A., 1996, Atom. Data Nucl. Data Tables, 64, 183  
 Bhatia A.K., Doschek G.A., Eissner W., 2002, Atom. Data Nucl. Data Tables, 82, 211  
 Brickhouse N.S., Raymond J.C., Smith B.W., 1995, ApJS, 97, 551  
 Brosius J.W., Davila J.M., Thomas R.J., Monsignori-Fossi B.C., 1996, ApJS, 106, 143  
 Brosius J.W., Davila J.M., Thomas R.J., 1998, ApJS, 119, 255  
 Brosius J.W., Thomas R.J., Davila J.M., Landi E., 2000, ApJ, 543, 1016  
 Brosius J.W., Rabin D.M., Thomas R.J., Landi E., 2008, ApJ, 677, 781  
 Brown C.M., Feldman U., Seely J.F., Korendyke C.M., Hara H., 2008, ApJS, 176, 511  
 Bryans P., Landi E., Savin D.W., 2009, ApJ, 691, 1540  
 Culhane J.L. et al., 2007, Solar Phys., 243, 19  
 Dere K.P., 1978, ApJ, 221, 1062  
 Dere K.P., Landi E., Mason H.E., Monsignori-Fossi B.C., Young P.R., 1997, A&AS, 125, 149  
 Dere K.P., Landi E., Young P.R., Del Zanna G., Landini M., Mason H.E., 2009, A&A, 498, 915  
 Dufton P.L., 1977, Comput. Phys. Commun., 13, 25  
 Dufton P.L., Berrington K.A., Burke P.G., Kingston A.E., 1978, A&A, 62, 111  
 Gupta G.P., Tayal S.S., 1999, ApJ, 510, 1078  
 Jupén C., Isler R.C., Träbert E., 1993, MNRAS, 264, 627  
 Kastner S.O., Mason H.E., 1978, A&A, 67, 119  
 Keenan F.P., Aggarwal K.M., Ryans R.S.I., Milligan R.O., Bloomfield D.S., Brosius J.W., Davila J.M., Thomas R.J., 2005, ApJ, 624, 428  
 Keenan F.P., Jess D.B., Aggarwal K.M., Thomas R.J., Brosius J.W., Davila J.M., 2007, MNRAS, 376, 205  
 Keenan F.P., Jess D.B., Aggarwal K.M., Thomas R.J., Brosius J.W., Davila J.M., 2008, MNRAS, 389, 939  
 Laming J.M., Drake J.J., 1999, ApJ, 516, 324  
 Landi E., Feldman U., Dere K.P., 2002, ApJ, 574, 495  
 Landman D.A., 1980, ApJ, 240, 709  
 Mason H.E. 1975, MNRAS, 170, 651  
 Neupert W.M., Epstein G.L., Thomas R.J., Thompson W.T., 1992, Solar Phys., 137, 87  
 Puchnarewicz E.M., Mason K.O., Siemiginowska A., 1998, MNRAS, 293, L52  
 Shirai T., Funatake Y., Mori K., Sugar J., Wiese W.L., Nakai Y., 1990, J. Phys. Chem. Ref. Data, 19, 127  
 Thomas R.J., Neupert W.M., 1994, ApJS, 91, 461  
 Träbert E., 1998, MNRAS, 297, 399  
 Young P.R., Landi E., Thomas R.J., 1998, A&A, 329, 291

This paper has been typeset from a  $\text{\TeX}/\text{\LaTeX}$  file prepared by the author.